

References

1. Miller MA, et al. Sleep and Cognition. In: Idzikowski C, ed. Sleep and its Disorders Affect Society. IntechOpen; 2014:chap 1.
2. Choudhary AK, et al. Sleep restriction may lead to disruption in physiological attention and reaction time. *Sleep Sci.* 2016;9:207-211.
3. Wesnes KA, Macher J-P. Modafinil reverses the marked attentional deficits produced by acute sleep deprivation in healthy volunteers. *J Psychopharmacol.* 2004;18(suppl):A48.
4. Harsh JR, et al. The efficacy and safety of armodafinil as treatment for adults with excessive sleepiness associated with narcolepsy. *Curr Med Res Opin.* 2006;22(4):761-774.
5. Czeisler CA, et al. Armodafinil for Treatment of Excessive Sleepiness Associated With Shift Work Disorder: A Randomized Controlled Study. Poster presented at: AAPA 2010.
6. Hirshkowitz M, et al. Adjunct armodafinil improves wakefulness and memory in obstructive sleep apnea/hypopnea syndrome. *Respir Med.* 2007;101(3):616-627.
7. Roth T, et al. Armodafinil improves wakefulness and long-term episodic memory in nCPAP-adherent patients with excessive sleepiness associated with obstructive sleep apnea. *Sleep Breath.* 2008;12(1):53-62.
8. Otmani S, et al. Effects of prolonged-release melatonin and zolpidem on postural stability in older adults. *Hum Psychopharmacol.* 2012;27(3):270-276.
9. Wesnes K, Warburton DM. A comparison of temazepam and flurazepam in terms of sleep quality and residual changes in performance. *Neuropsychobiology.* 1984;11(4):255-259.
10. Deacon S, et al. Effect of short-term treatment with gaboxadol on sleep maintenance and initiation in patients with primary insomnia. *Sleep.* 2007;30(3):281-287.