

References

1. Armstrong AW, Parsi K, Schupp CW, Mease PJ, Duffin KC. Standardizing training for psoriasis measures: effectiveness of an online training video on Psoriasis Area and Severity Index assessment by physician and patient raters. *JAMA Dermatol.* 2013;149(5):577-582.
2. Božek A, Reich A. Assessment of intra-and inter-rater reliability of three methods for measuring atopic dermatitis severity: EASI, objective SCORAD, and IGA. *Dermatology.* 2017;233(1):16-22.
3. Everhart A, Kalpadakis-Smith A. Properly training physicians in dermatology trials. *Appl Clin Trials.* 2021. <https://www.appliedclinicaltrials.com/view/properly-training-physicians-in-dermatology-trials>
4. Silverberg JI, Simpson B, Abuabara K, et al; TARGET-DERM Investigators. Prevalence and burden of atopic dermatitis involving the head, neck, face, and hand: a cross sectional study from the TARGET-DERM AD cohort. *J Am Acad Dermatol.* 2023;89(3):519-528.
5. Williams HC, Schmitt J, Thomas KS, et al; HOME Initiative. The HOME Core outcome set for clinical trials of atopic dermatitis. *J Allergy Clin Immunol.* 2022;149(6):1899-1911.
6. Youn SW, Choi CW, Kim BR, Chae JB. Reduction of inter-rater and intra-rater variability in psoriasis area and severity index assessment by photographic training. *Ann Dermatol.* 2015;27(5):557.